

AVANOS

ON-Q* DELIVERS

One partner. One mission.
Unmatched service and support.



Proven opioid-sparing
perioperative pain
management¹⁻⁴

Widest selection
of elastomeric *and*
electronic pumps

Customizable
to meet every
patient's needs^{2,5-9}

Δ SHARED PARTNERSHIP IN THE FIGHT AGAINST OPIOID DEPENDENCY

ON-Q* + You

Dependency can begin **within just 3** days of initial opioid use.¹⁰



As many as **1 in 5 patients** becomes a routine opioid user after **10 days** of narcotic analgesia.¹⁰



EVEN A 1-DAY OPIOID PRESCRIPTION MAY POSE A 6% RISK OF LONG-TERM USE.¹⁰

NOW OFFERING EVEN MORE OPTIONS FOR OPIOID-SPARING PAIN RELIEF

ON-Q* + ambIT



ELASTOMERIC PAIN PUMPS

- Up to **5 days** of targeted pain relief ^{11,12}
- Allows for **adjustable flow** during therapy
- **Bolus option** for breakthrough pain
- **On/Off** capability throughout therapy



ELECTRONIC PAIN PUMPS

- Up to **1000ml** of local anesthetic¹³
- **Programmable intermittent bolus** option
- Bolus button with **safety lockout**
- Audible/visual **infusion confirmation**

UNMATCHED SERVICE AND SUPPORT. ALL DAY. EVERY DAY.

ON-Q* + ambIT



PATIENT AND CLINICIAN STAFF SUPPORT

24/7 hotline staffed by nurses, training of your existing and new staff, **ongoing clinician in-servicing**, pain catheter insertion training for continuous blocks and surgical site placement, **patient at-home** and clinician guides

EXTENSIVE LOCAL SUPPORT

Coordinated ON-Q* representatives and clinical experts to **facilitate uninterrupted personalized coverage and continuing success**

ONGOING TRAINING

Helping surgeons, anesthesiologists, and staff **stay current with techniques** and **keep workflows efficient**

THE SIMPLE CHOICE FOR CUSTOMIZED OPIOID-SPARING PROTOCOLS

ON-Q* + ambIT



FLEXIBILITY

- **More choices:** elastomeric and electronic pain pumps offering adjustable flow, on-demand and intermittent bolus options
- **Opioid-free pain relief:** integral to multimodal plans^{1-4,12}
- **Customizable controls:** titratability, patient bolus delivery, pre-set protocols, programmable intermittent bolus, and more



IMPROVED OUTCOMES

- Significantly **reduced opioid use**¹⁻⁴
- **Shortened hospital stays** by 1.1 days^{2,5,9,14-22}
- Effective, long-lasting pain relief from **hospital to post-discharge**^{2,5-9}
- Patients up to **3x likelier** to report high satisfaction vs opioids^{9,14,15,23-25}



INCREASED EFFICIENCIES

- **Single-source partner** for all of your ambulatory pain pump needs
- Easy pump setup that **does not prolong procedure time**
- **Cost-effectiveness** through shorter stays, fewer complications, lower readmission rates^{1,2,5,8,9,14-23,26-29}

AVANOS

**CHOOSE THE ON-Q*
AMBULATORY PAIN
MANAGEMENT
PORTFOLIO.**

**NOW INCLUDING
ambIT®.**

Your single source for customized,
opioid-sparing perioperative pain
management.



REFERENCES: **1.** Benyamin R, Trescot AM, Datta S, et al. Opioid complications and side effects. *Pain Physician*. 2008;11:s105-s120. **2.** Forastiere E, Sofra M, Giannarelli D, Fabrizi L, Simone G. Effectiveness of continuous wound infusion of 0.5% ropivacaine by ON-Q pain relief system for postoperative pain management after open nephrectomy. *Br J Anaesth*. 2008;101(6):841-847. **3.** Parvizi J, Miller AG, Gandhi K. Multimodal pain management after total joint arthroplasty. *J Bone Joint Surg Am*. 2011;93(11):1075-1184. **4.** Webb C, Mariano E. Best multimodal analgesic protocol for total knee arthroplasty. *Pain Management*. 2015;5(3):185-196. **5.** Auyong DB, Allen CJ, Pahang JA, Clabeaux JJ, MacDonald KM, Hanson NA. Reduced length of hospitalization in primary total knee arthroplasty patients using an updated enhanced recovery after orthopedic surgery (ERAS) pathway. *J Arthroplasty*. 2015;30(10):1705-1709. **6.** Bingham AE, Fu R, Horn JL, Abrahams MS. Continuous peripheral nerve block compared with single-injection peripheral nerve block: a systematic review and meta analysis of randomized controlled trials. *Reg Anesth Pain Med*. 2012;37(6):583-594. **7.** Dine A. Evidence based outcomes review. Continuing review and evaluation for I-Flow Corporation. July 2012. **8.** Goyal N, McKenzie J, Sharkey PF, Parvizi J, Hozack WJ, Austin MS. The 2012 Chitranjan Ranawat award: intraarticular analgesia after TKA reduces pain: a randomized, double-blinded, placebo-controlled, prospective study. *Clin Orthop Relat Res*. 2013;471(1):64-75. **9.** Liu SS, Richman JM, Thirlby RC, Wu CL. Efficacy of continuous wound catheters delivering local anesthetic for postoperative analgesia: a quantitative and qualitative systematic review of randomized controlled trials. *J Am Coll Surg*. 2006;203(6):914-932. **10.** Centers for Disease Control and Prevention (CDC). Characteristics of initial prescription episodes and likelihood of long-term opioid use—United States, 2006-2015. <https://www.cdc.gov/mmwr/volumes/66/wr/mm6610a1.htm>. Accessed February 7, 2018. **11.** Abdallah FW, Halpern SH, Aoyama K, Brull R. Will the real benefits of single-shot interscalene block please stand up? A systematic review and meta-analysis. *Anesth Analg*. 2015. 120(5):1114-1129. **12.** IFU - The ON-Q* Pain Relief System. Instructions for use. **13.** ambit® Pump Clinician Manual. Summit Medical Products. May 2019. **14.** Beaussier M, El'Ayoubi H, Schiffer E, et al. Continuous preperitoneal infusion of ropivacaine provides effective analgesia and accelerates recovery after colorectal surgery. *Anesthesiology*. 2007;107(3):461-468. **15.** Bianconi M, Ferraro L, Traina GC, et al. Pharmacokinetics and efficacy of ropivacaine continuous wound instillation after joint replacement surgery. *Br J Anaesth*. 2003;91(6):830-835. **16.** Capdevila X, Barthelet Y, Ryckwaert Y, Rubenovitch J, d'Athis F. Effects of perioperative analgesic technique on the surgical outcome and duration of rehabilitation after major knee surgery. *Anesthesiology*. 1999;91(1):8-15. **17.** Carli F, Clemente A, Asenjo JF, et al. Analgesia and functional outcome after total knee arthroplasty: periarticular infiltration vs continuous femoral nerve block. *Br J Anaesth*. 2010;105(2):185-195. **18.** Dowling R, Thielmeier K, Ghaly A, Barber D, Boice T, Dine A. Improved pain control after cardiac surgery: results of a randomized, double-blind, clinical trial. *J Thorac Cardiovasc Surg*. 2003;126(5):1271-1278. **19.** Ilfeld BM, Ball ST, Gearen PF, et al. Ambulatory continuous posterior lumbar plexus nerve blocks after hip arthroplasty: a dual-center, randomized, triple-masked, placebo controlled trial. *Anesthesiology*. 2008;109(3):491-501. **20.** Mudumbai SC, Kim TE, Howard SK, et al. Continuous adductor canal blocks are superior to continuous femoral nerve blocks in promoting early ambulation after TKA. *Clin Orthop Relat Res*. 2014;472(5):1377-1383. **21.** White PF, Rawal S, Latham P, Markowitz S, et al. Use of a continuous local anesthetic infusion for pain management after median sternotomy. *Anesthesiology*. 2003;99(4):918-923. **22.** Mangram AJ, Oguntodu OF, Hollingworth AK, Prokuski L, Steinstra A, Collins M, et al. Geriatric trauma G-60 falls with hip fractures: A pilot study of acute pain management using femoral nerve fascia iliac blocks. *Journal of Trauma and Acute Care Surgery*. 2015Dec;79(6):1067-72. **23.** Heller L, Kowalski AM, Wei C, Butler CE. Prospective, randomized, double-blind trial of local anesthetic infusion and intravenous narcotic patient-controlled anesthesia pump for pain management after free TRAM flap breast reconstruction. *Plast Reconstr Surg*. 2008;122(4):1010-1018. **24.** Husain A, Lee GC. Establishing realistic patient expectations following total knee arthroplasty. *J Am Acad Orthop Surg*. 2015;23(12):707-713. **25.** Kean J, Wigderowitz CA, Coventry DM. Continuous interscalene infusion and single injection using levobupivacaine for analgesia after surgery of the shoulder: a double blind, randomized controlled trial. *J Bone Joint Surg [Br]*. 2006;88-B:1173-1177. **26.** American Academy of Orthopaedic Surgeons Information statement. Opioid use, misuse, and abuse in orthopaedic practice. https://www.aaos.org/uploadedFiles/PreProduction/About/Opinion_Statements/advistmt/1045%20Opioid%20Use,%20Misuse,%20and%20Abuse%20in%20Practice.pdf. Published October 2015. Accessed February 14, 2018. **27.** Aguirre J, Del Moral A, Cobo I, Borgeat A, Blumenthal S. The role of continuous peripheral nerve blocks. *Anesthesiol Res Pract*. 2012;2012:1-20. **28.** Chelly JE, Ghisi D, Fanelli A. Continuous peripheral nerve blocks in acute pain management. *Br J Anaesth*. 2010;105(suppl 1):86-96. **29.** The Joint Commission. Safe use of opioids in hospitals. *Sentinel Event Alert*. Issue 49, August 8, 2012.

For more information, please visit
avanos.com
1-844-4AVANOS (1-844-428-2667)

There are inherent risks in all medical devices. Please refer to the product labeling for Indications, Cautions, and Warnings and Contraindications. Failure to follow the product labeling could directly impact patient safety. Physician is responsible for prescribing and administering medications per instructions provided by the drug manufacturer. Refer to www.avanos.com for product safety Technical Bulletins.

*Registered Trademark or Trademark of Avanos Medical, Inc., or its affiliates.
©2019 AVNS. All rights reserved. COPY-XXXX 11/19

AVANOS