

TECHNICAL BULLETIN

WHAT WE KNOW ABOUT CHONDROLYSIS

A number of papers have been published on the subject of chondrolysis. Early papers tended to focus on thermal capsulorrhaphy. In a paper published in the spring of 2004, Petty et al¹ reported on three case histories and offered several possible causes of the condition. In July 2006 Gomoll et al² reported on a possible connection of chondrolysis to continuous intra-articular infusion of bupivacaine based on studies performed on rabbit shoulders. In August of 2007, Hansen et al³ published a paper based on a retrospective review of 177 shoulder arthroscopies of 152 patients of the senior author. Hansen described postarthroscopic glenohumeral chondrolysis (PAGCL) in 12 shoulders of 10 patients that they suggest are related to the continuous intra-articular use of bupivacaine with epinephrine. It is noteworthy that Hansen reported that 104 of the patients had bupivacaine administered extra-articularly with no incidence of PAGCL in this group. Hansen summarized other possible causes of reported chondrolysis cases that have been identified in recent years by a variety of investigators. Some of the possible causes as presented in various papers, up to and including the most recent Hansen paper, are as follows:

- Thermal capsulorrhaphy^{4, 5}
- Radiofrequency treatment⁶
- Prominent hardware⁷
- Gentian Violet⁸
- Chlorhexidine⁹
- Methylmethacrylate¹⁰
- Intra-articular administration of bupivacaine with epinephrine³
- Intra-articular administration of bupivacaine^{2, 11}

Because of the vasoconstrictive properties of epinephrine, in 2003 to be proactive, I-Flow* (currently, Avanos Medical) incorporated the following warning regarding the use of epinephrine in its devices:

VASOCONSTRICTORS SUCH AS EPINEPHRINE ARE NOT RECOMMENDED FOR CONTINUOUS INFUSIONS.

In late 2006, in light of the possible connection between the use of bupivacaine in the intra-articular space and the onset of chondrolysis as suggested by the Gomoll paper, and in an abundance of caution, I-Flow* (currently, Avanos Medical) modified the ON-Q* DFU by adding the following warning:

Avoid placing the catheter in joint spaces. Although there is no definitive established causal relationship, some literature has shown a possible association between continuous intra-articular infusions (particularly with bupivacaine) and the subsequent development of chondrolysis.

ON-Q* Pain Management Systems are indicated for intra-operative applications, among other applications. The literature raises concerns regarding intra-articular use. The attached bibliography is for your reference. **As always, the decision on how to treat the patient and what medications to administer belongs exclusively to the physician.**

REFERENCES:

1. Petty DH, Jazrawi LM, Estrada LS, Andrew JR. Glenohumeral chondrolysis after shoulder arthroscopy: case reports and review of the literature. *Am J Sports Med* 2004;32(2):509-15.
2. Gomoll AH, Kang RW, Williams JM, Bach BR, Cole BJ. Chondrolysis after continuous intra-articular bupivacaine infusion: an experimental model investigating chondrotoxicity in the rabbit shoulder. *Arthroscopy* 2006;22(8):813-19.
3. Hansen BP, Beck CL, Beck EP, Townsley RW. Postarthroscopic glenohumeral chondrolysis. *Am J Sports Med*. 2007; epub 2007 July 3 :1-7.
4. Levine WN, Clark AM, D'Alessandro FD, Yamaguchi K. Chondrolysis following arthroscopic thermal capsulorrhaphy to treat shoulder instability. *J Bone Joint Surg* 2005;87A(3):616-21.
5. Ciccione WJ, Weinstein DM, Elias JJ. Glenohumeral chondrolysis following thermal capsulorrhaphy. *Orthopedics* 2007;30(2):158-60.
6. Jerosch J, Aldawoudy AM. Chondrolysis of the glenohumeral joint following arthroscopic capsular release for adhesive capsulitis: a case report. *Knee Surg Sports Traumatol Arthrosc* 2007;15:292-4.
7. Sternlicht AL, Ehrlich MG, Armstrong AL, Zaleske DJ. Role of pin protrusion in the etiology of chondrolysis: a surgical model with radiographic, histologic, and biomechanical analysis. *J Pediatr Orthop* 1992;12(4):428-33.
8. Shibata Y, Midorikawa K, Koga T, Honjo N, Naito M. Chondrolysis of the glenohumeral joint following a color test using gentian violet. *International Orthopaedics* 2001;25:401-403.
9. van Huyssten AL, Bracey DJ. Chlorhexidine and chondrolysis in the knee. *J Bone Joint Surg* 1999;81-B:995-6.
10. Leclair A, Gangi A, Lacaze F. Rapid chondrolysis after an intra-articular leak of bone cement in treatment of a benign acetabular subchondral cyst: an unusual complication of percutaneous injection of acrylic cement. *Skeletal Radiol*. 2000;29(5):275-78.
11. Chu CR, Izzo NJ, Papas NE, Fu F. In vitro exposure to 0.5% bupivacaine is cytotoxic to bovine articular chondrocytes. *Arthroscopy* 2006;22(7):693-9.

There are inherent risks in all medical devices. Please refer to the product labeling for **Indications, Cautions, Warnings** and **Contraindications**. Failure to follow the product labeling could directly impact patient safety. Physician is responsible for prescribing and administering medications per instructions provided by the drug manufacturer. Refer to www.avanospainmanagement.com for additional product safety Technical Bulletins.

Please contact the Clinical Services Department at **800-444-2728** or **949-923-2400** if you have any questions regarding this information.

Page 2 of 2

AVANOS

[avanospainmanagement.com](http://www.avanospainmanagement.com)



Rx only.

*Registered Trademark or Trademark of Avanos Medical, Inc., or its affiliates.

© 2018 AVNS. All rights reserved.

MK-00015 rev2 07/2015